


Marine Fossilien aus der Böhlen-Formation (Unteroligozän) des Leipziger Südraumes


Gastropoda (Schnecken)

Fissurellidae




Emarginula nystiana BOSQUET, 1851

Trochidae




Solariella suturalis (PHILIPPI, 1843)



Margarites margaritula (SANDBERGER, 1843)

Capulidae




Calyptreaa striatella NYST, 1845

Xenophoridae




Tugurium scrutarium (PHILIPPI, 1843)

Naticidae




Lunatia dilatata (PHILIPPI, 1843)



Polinices? hantoniensis (PILKINGTON, 1804)

Cassidae



Phalium rondeleti (Solander, 1766)



Galeodea nodosa (SOLANDER, 1766)

Cymatiidae




Charonia flandrica (DE KONINCK, 1837)

Aporrhaidae



Drepanocheilus speciosus (SCHLOTHEIM, 1820)

Muricidae



Trophonopsis pauwelsii (DE KONINCK, 1837)




Boreotrophon capito (PHILIPPI, 1843)

Nassariidae




Keepingia gossardi (NYST, 1836)

Buccinidae



Scalaspira multisulcata (NYST, 1845)

Scalaspira waeli (BEYRICH, 1856)



Scalaspira convexa (SANDBERGER, 1860)

Scalaspira villana (PHILIPPI, 1847)


Turbinellidae



Exilia elatior (BEYRICH, 1848)

Streptodictyon sowerbyi (NYST, 1836)

Fascioliariidae




Volutidae



Scaphella siemsseni (BOLL, 1851)

Cancellariidae




Cancellaria evulsa (SOLANDER, 1766)

Turridae



Bathytoma leunisi (PHILIPPI, 1843)


Bathytoma subdenticulata (MÜNSTER, 1835)



Acamplogenotia morreni (DE KONINCK, 1837)

Fusiturris selysii (DE KONINCK, 1837)

Conidae




Conus symmetricus (SANDBERGER, 1859)

Architectonicidae




Architectonica dumonti (NYST, 1845)

Actaeonidae



Tornatellaea simulata (SOLANDER, 1766)

Cavoliniidae



Creseis maxima (LUDWIG, 1864)